

Frontiers in Immunology, 2017, vol.8, NNOV

## Immunoglobulins G from sera of amyotrophic lateral sclerosis patients induce oxidative stress and upregulation of antioxidative system in BV-2 microglial cell line

Milošević M., Milicević K., Božić I., Lavrnja I., Stevanović I., Bijelić D., Dubaić M., Ivković I., Stević Z., Giniatullin R., Andjus P.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

### Abstract

© 2017 Milošević, Milicević, Božić, Lavrnja, Stevanović, Bijelić, Dubaić, Živković, Stević, Giniatullin and Andjus. Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disorder with a very fast progression, no diagnostic tool for the presymptomatic phase, and still no effective treatment of the disease. Although ALS affects motor neurons, the overall pathophysiological condition points out to the non-cell autonomous mechanisms, where astrocytes and microglia play crucial roles in the disease progression. We have already shown that IgG from sera of ALS patients (ALS IgG) induce calcium transients and an increase in the mobility of acidic vesicles in cultured rat astrocytes. Having in mind the role of microglia in neurodegeneration, and a well-documented fact that oxidative stress is one of the many components contributing to the disease, we decided to examine the effect of ALS IgG on activation, oxidative stress and antioxidative system of BV-2 microglia, and to evaluate their acute effect on cytosolic peroxide, pH, and on reactive oxygen species (ROS) generation. All tested ALS IgGs (compared to control IgG) induced oxidative stress (rise in nitric oxide and the index of lipid peroxidation) followed by release of TNF- $\alpha$  and higher antioxidative defense (elevation of Mn- and CuZn-superoxide dismutase, catalase, and glutathione reductase with a decrease of glutathione peroxidase and glutathione) after 24 h treatment. Both ALS IgG and control IgG showed same localization on the membrane of BV-2 cells following 24 h treatment. Cytosolic peroxide and pH alteration were evaluated with fluorescent probes HyPer and SypHer, respectively, having in mind that HyPer also reacts to pH changes. Out of 11 tested IgGs from ALS patients, 4 induced slow exponential rise of HyPer signal, with maximal normalized fluorescence in the range 0.2-0.5, also inducing similar increase of SypHer intensity, but of a lower amplitude. None of the control IgGs induced changes with neither of the indicators. Acute ROS generation was detected in one out of three tested ALS samples with carboxy-H2DCFDA. The observed phenomena demonstrate the potential role of inflammatory humoral factors, IgGs, as potential triggers of the activation in microglia, known to occur in later stages of ALS. Therefore, revealing the ALS IgG signaling cascade in microglial cells could offer a valuable molecular biomarker and/or a potential therapeutic target.

<http://dx.doi.org/10.3389/fimmu.2017.01619>

## Keywords

Amyotrophic lateral sclerosis, Antioxidative system, BV-2 microglia, HyPer, Immunoglobulin G, Oxidative stress, SypHer

## References

- [1] Kiernan MC, Vucic S, Cheah BC, Turner MR, Eisen A, Hardiman O, et al. Amyotrophic lateral sclerosis. *Lancet* (2011) 377:942-55. doi:10.1016/S0140-6736(10)61156-7
- [2] Alexianu ME, Kozovska M, Appel SH. Immune reactivity in a mouse model of familial ALS correlates with disease progression. *Neurology* (2001) 57:1282-9. doi:10.1212/WNL.57.7.1282
- [3] Engelhardt JI, Appel SH. IgG reactivity in the spinal cord and motor cortex in amyotrophic lateral sclerosis. *Arch Neurol* (1990) 47:1210-6. doi:10.1001/archneur.1990.00530110068019
- [4] Kawamata T, Akiyama H, Yamada T, McGeer PL. Immunologic reactions in amyotrophic lateral sclerosis brain and spinal cord tissue. *Am J Pathol* (1992) 140:691-707
- [5] McGeer PL, McGeer EG. Inflammatory processes in amyotrophic lateral sclerosis. *Muscle Nerve* (2002) 26:459-70. doi:10.1002/mus.10191
- [6] Turner MR, Cagnin A, Turkheimer FE, Miller CC, Shaw CE, Brooks DJ, et al. Evidence of widespread cerebral microglial activation in amyotrophic lateral sclerosis: an [<sup>11</sup>C](R)-PK11195 positron emission tomography study. *Neurobiol Dis* (2004) 15:601-9. doi:10.1016/j.nbd.2003.12.012
- [7] Clement AM, Nguyen MD, Roberts EA, Garcia ML, Boillee S, Rule M, et al. Wild-type nonneuronal cells extend survival of SOD1 mutant motor neurons in ALS mice. *Science* (2003) 302:113-7. doi:10.1126/science.1086071
- [8] Frakes AE, Ferraiuolo L, Haidet-Phillips AM, Schmelzer L, Braun L, Miranda CJ, et al. Microglia induce motor neuron death via the classical NF-kappaB pathway in amyotrophic lateral sclerosis. *Neuron* (2014) 81:1009-23. doi:10.1016/j.neuron.2014.01.013
- [9] Kabba JA, Xu Y, Christian H, Ruan W, Chenai K, Xiang Y, et al. Microglia: housekeeper of the central nervous system. *Cell Mol Neurobiol* (2017). doi:10.1007/s10571-017-0504-2
- [10] Waisman A, Ginhoux F, Greter M, Bruttger J. Homeostasis of microglia in the adult brain: review of novel microglia depletion systems. *Trends Immunol* (2015) 36:625-36. doi:10.1016/j.it.2015.08.005
- [11] Wlodarczyk A, Cedile O, Jensen KN, Jasson A, Mony JT, Khorooshi R, et al. Pathologic and protective roles for microglial subsets and bone marrow-and blood-derived myeloid cells in central nervous system inflammation. *Front Immunol* (2015) 6:463. doi:10.3389/fimmu.2015.00463
- [12] Ginhoux F, Lim S, Hoeffel G, Low D, Huber T. Origin and differentiation of microglia. *Front Cell Neurosci* (2013) 7:45. doi:10.3389/fncel.2013.00045
- [13] Hanisch UK, Kettenmann H. Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat Neurosci* (2007) 10:1387-94. doi:10.1038/nn1997
- [14] Henkel JS, Beers DR, Zhao W, Appel SH. Microglia in ALS: the good, the bad, and the resting. *J Neuroimmune Pharmacol* (2009) 4:389-98. doi:10.1007/s11481-009-9171-5
- [15] Kreutzberg GW. Microglia: a sensor for pathological events in the CNS. *Trends Neurosci* (1996) 19:312-8. doi:10.1016/0166-2236(96)10049-7
- [16] Ransohoff RM, Perry VH. Microglial physiology: unique stimuli, specialized responses. *Annu Rev Immunol* (2009) 27:119-45. doi:10.1146/annurev.immunol.021908.132528
- [17] Boillee S, Yamanaka K, Lobsiger CS, Copeland NG, Jenkins NA, Kassiotis G, et al. Onset and progression in inherited ALS determined by motor neurons and microglia. *Science* (2006) 312:1389-92. doi:10.1126/science.1123511
- [18] Yamanaka K, Chun SJ, Boillee S, Fujimori-Tonou N, Yamashita H, Gutmann DH, et al. Astrocytes as determinants of disease progression in inherited amyotrophic lateral sclerosis. *Nat Neurosci* (2008) 11:251-3. doi:10.1038/nn2047
- [19] Bataveljic D, Milosevic M, Radenovic L, Andjus P. Novel molecular biomarkers at the blood-brain barrier in ALS. *Biomed Res Int* (2014) 2014:907545. doi:10.1155/2014/907545
- [20] Stenovec M, Milosevic M, Petrusic V, Potokar M, Stevic Z, Prebil M, et al. Amyotrophic lateral sclerosis immunoglobulins G enhance the mobility of Lysotracker-labelled vesicles in cultured rat astrocytes. *Acta Physiol* (2011) 203:457-71. doi:10.1111/j.1748-1716.2011.02337.x
- [21] Milosevic M, Stenovec M, Kreft M, Petrusic V, Stevic Z, Trkov S, et al. Immunoglobulins G from patients with sporadic amyotrophic lateral sclerosis affects cytosolic Ca<sup>2+</sup> homeostasis in cultured rat astrocytes. *Cell Calcium* (2013) 54:17-25. doi:10.1016/j.ceca.2013.03.005

- [22] Mishra PS, Dhull DK, Nalini A, Vijayalakshmi K, Sathyaprabha TN, Alladi PA, et al. Astroglia acquires a toxic neuroinflammatory role in response to the cerebrospinal fluid from amyotrophic lateral sclerosis patients. *J Neuroinflammation* (2016) 13:212. doi:10.1186/s12974-016-0698-0
- [23] Sharma A, Varghese AM, Vijayalakshmi K, Sumitha R, Prasanna VK, Shruthi S, et al. Cerebrospinal fluid from sporadic amyotrophic lateral sclerosis patients induces mitochondrial and lysosomal dysfunction. *Neurochem Res* (2016) 41:965-84. doi:10.1007/s11064-015-1779-7
- [24] Obal I, Klausz G, Mandi Y, Deli M, Siklos L, Engelhardt JI. Intraperitoneally administered IgG from patients with amyotrophic lateral sclerosis or from an immune-mediated goat model increase the levels of TNF-alpha, IL-6, and IL-10 in the spinal cord and serum of mice. *J Neuroinflammation* (2016) 13:121. doi:10.1186/s12974-01-0586-7
- [25] Pullen AH, Demestre M, Howard RS, Orrell RW. Passive transfer of purified IgG from patients with amyotrophic lateral sclerosis to mice results in degeneration of motor neurons accompanied by Ca<sup>2+</sup> enhancement. *Acta Neuropathol* (2004) 107:35-46. doi:10.1007/s00401-003-0777-z
- [26] Garbuzova-Davis S, Sanberg PR. Blood-CNS barrier impairment in ALS patients versus an animal model. *Front Cell Neurosci* (2014) 8:21. doi:10.3389/fncel.2014.00021
- [27] An T, Shi P, Duan W, Zhang S, Yuan P, Li Z, et al. Oxidative stress and autophagic alteration in brainstem of SOD1-G93A mouse model of ALS. *Mol Neurobiol* (2014) 49:1435-48. doi:10.1007/s12035-013-8623-3
- [28] Barber SC, Shaw PJ. Oxidative stress in ALS: key role in motor neuron injury and therapeutic target. *Free Radic Biol Med* (2010) 48:629-41. doi:10.1016/j.freeradbiomed.2009.11.018
- [29] Bergeron C. Oxidative stress: its role in the pathogenesis of amyotrophic lateral sclerosis. *J Neurol Sci* (1995) 129:81-4. doi:10.1016/0022-510X(95)00071-9
- [30] Ferrante RJ, Browne SE, Shinobu LA, Bowling AC, Baik MJ, MacGarvey U, et al. Evidence of increased oxidative damage in both sporadic and familial amyotrophic lateral sclerosis. *J Neurochem* (1997) 69:2064-74. doi:10.1046/j.1471-4159.1997.69052064.x
- [31] Ikawa M, Okazawa H, Tsujikawa T, Matsunaga A, Yamamura O, Mori T, et al. Increased oxidative stress is related to disease severity in the ALS motor cortex: a PET study. *Neurology* (2015) 84:2033-9. doi:10.1212/WNL.0000000000001588
- [32] Stamenkovic S, Pavicevic A, Mojovic M, Popovic-Bijelic A, Selakovic V, Andjus P, et al. In vivo EPR pharmacokinetic evaluation of the redox status and the blood brain barrier permeability in the SOD1G93A ALS rat model. *Free Radic Biol Med* (2017) 108:258-69. doi:10.1016/j.freeradbiomed.2017.03.034
- [33] Brooks BR, Miller RG, Swash M, Munsat TL; World Federation of Neurology Research Group on Motor Neuron Diseases. El Escorial revisited: revised criteria for the diagnosis of amyotrophic lateral sclerosis. *Amyotroph Lateral Scler Other Motor Neuron Disord* (2000) 1:293-9
- [34] Blasi E, Barluzzi R, Bocchini V, Mazzolla R, Bistoni F. Immortalization of murine microglial cells by a v-raf/v-myc carrying retrovirus. *J Neuroimmunol* (1990) 27:229-37. doi:10.1016/0165-5728(90)90073-V
- [35] Andjus PR, Stevic-Marinkovic Z, Cherubini E. Immunoglobulins from motoneurone disease patients enhance glutamate release from rat hippocampal neurones in culture. *J Physiol* (1997) 504:103-12. doi:10.1111/j.1469-7793.1997.103bf.x
- [36] Navarro-Gonzalez JA, Garcia-Benayas C, Arenas J. Semiautomated measurement of nitrate in biological fluids. *Clin Chem* (1998) 44:679-81
- [37] Villacara A, Kumami K, Yamamoto T, Mrsulja BB, Spatz M. Ischemic modification of cerebrocortical membranes: 5-hydroxytryptamine receptors, fluidity, and inducible in vitro lipid peroxidation. *J Neurochem* (1989) 53:595-601. doi:10.1111/j.1471-4159.1989.tb07375.x
- [38] Sun M, Zigman S. An improved spectrophotometric assay for superoxide dismutase based on epinephrine autooxidation. *Anal Biochem* (1978) 90:81-9. doi:10.1016/0003-2697(78)90010-6
- [39] Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem* (1951) 193:265-75
- [40] Goth L. A simple method for determination of serum catalase activity and revision of reference range. *Clin Chim Acta* (1991) 196:143-51. doi:10.1016/0009-8981(91)90067-M
- [41] Djukic MM, Jovanovic MD, Ninkovic M, Stevanovic I, Ilic K, Curcic M, et al. Protective role of glutathione reductase in paraquat induced neurotoxicity. *Chem Biol Interact* (2012) 199:74-86. doi:10.1016/j.cbi.2012.05.008
- [42] Maral J, Puget K, Michelson AM. Comparative study of superoxide dismutase, catalase and glutathione peroxidase levels in erythrocytes of different animals. *Biochem Biophys Res Commun* (1977) 77:1525-35. doi:10.1016/S0006-291X(77)80151-4
- [43] Freifelder D. *Physical Biochemistry: Applications to Biochemistry and Molecular Biology*. San Francisco: W. H. Freeman (1976)
- [44] Anderson M, Greenwald R. *Handbook of Methods for Oxygen Radical Research*. Florida: CRC Press Inc (1985). p. 317-23

- [45] Markvicheva KN, Bilan DS, Mishina NM, Gorokhovatsky AY, Vinokurov LM, Lukyanov S, et al. A genetically encoded sensor for H<sub>2</sub>O<sub>2</sub> with expanded dynamic range. *Bioorg Med Chem* (2011) 19:1079-84. doi:10.1016/j.bmc.2010.07.014
- [46] Belousov VV, Fradkov AF, Lukyanov KA, Staroverov DB, Shakhbazov KS, Terskikh AV, et al. Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nat Methods* (2006) 3:281-6. doi:10.1038/nmeth866
- [47] Rojo AI, McBean G, Cindric M, Egea J, Lopez MG, Rada P, et al. Redox control of microglial function: molecular mechanisms and functional significance. *Antioxid Redox Signal* (2014) 21:1766-801. doi:10.1089/ars.2013.5745
- [48] Liu Y, Kintner DB, Chanana V, Algharabli J, Chen X, Gao Y, et al. Activation of microglia depends on Na<sup>+</sup>/H<sup>+</sup> exchange-mediated H<sup>+</sup> homeostasis. *J Neurosci* (2010) 30:15210-20. doi:10.1523/JNEUROSCI.3950-10.2010
- [49] Beers DR, Henkel JS, Zhao W, Wang J, Appel SH. CD4<sup>+</sup> T cells support glial neuroprotection, slow disease progression, and modify glial morphology in an animal model of inherited ALS. *Proc Natl Acad Sci U S A* (2008) 105:15558-63. doi:10.1073/pnas.0807419105
- [50] Sta M, Sylva-Steenland RM, Casula M, de Jong JM, Troost D, Aronica E, et al. Innate and adaptive immunity in amyotrophic lateral sclerosis: evidence of complement activation. *Neurobiol Dis* (2011) 42:211-20. doi:10.1016/j.nbd.2011.01.002
- [51] Appel SH, Engelhardt JI, Garcia J, Stefani E. Immunoglobulins from animal models of motor neuron disease and from human amyotrophic lateral sclerosis patients passively transfer physiological abnormalities to the neuromuscular junction. *Proc Natl Acad Sci U S A* (1991) 88:647-51. doi:10.1073/pnas.88.2.647
- [52] Demestre M, Pullen A, Orrell RW, Orth M. ALS-IgG-induced selective motor neurone apoptosis in rat mixed primary spinal cord cultures. *J Neurochem* (2005) 94:268-75. doi:10.1111/j.1471-4159.2005.03184.x
- [53] Andjus PR, Khiroug L, Nistri A, Cherubini E. ALS IgGs suppress [Ca<sup>2+</sup>]<sub>i</sub> rise through P/Q-type calcium channels in central neurones in culture. *Neuroreport* (1996) 7:1914-6. doi:10.1097/00001756-199608120-00008
- [54] Engelhardt JI, Siklos L, Komuves L, Smith RG, Appel SH. Antibodies to calcium channels from ALS patients passively transferred to mice selectively increase intracellular calcium and induce ultrastructural changes in motoneurons. *Synapse* (1995) 20:185-99. doi:10.1002/syn.890200302
- [55] Llinas R, Sugimori M, Cherksey BD, Smith RG, Delbono O, Stefani E, et al. IgG from amyotrophic lateral sclerosis patients increases current through P-type calcium channels in mammalian cerebellar Purkinje cells and in isolated channel protein in lipid bilayer. *Proc Natl Acad Sci U S A* (1993) 90:11743-7. doi:10.1073/pnas.90.24.11743
- [56] Pagani MR, Reisin RC, Uchitel OD. Calcium signaling pathways mediating synaptic potentiation triggered by amyotrophic lateral sclerosis IgG in motor nerve terminals. *J Neurosci* (2006) 26:2661-72. doi:10.1523/JNEUROSCI.4394-05.2006
- [57] Engelhardt JI, Siklos L, Appel SH. Altered calcium homeostasis and ultrastructure in motoneurons of mice caused by passively transferred anti-motoneuronal IgG. *J Neuropathol Exp Neurol* (1997) 56:21-39. doi:10.1097/00005072-199701000-00003
- [58] Edri-Brami M, Rosental B, Hayoun D, Welt M, Rosen H, Wirguin I, et al. Glycans in sera of amyotrophic lateral sclerosis patients and their role in killing neuronal cells. *PLoS One* (2012) 7(5):e35772. doi:10.1371/journal.pone.0035772
- [59] Edri-Brami M, Sharoni H, Hayoun D, Skutelsky L, Nemirovsky A, Porgador A, et al. Development of stage-dependent glycans on the Fc domains of IgG antibodies of ALS animals. *Exp Neurol* (2015) 267:95-106. doi:10.1016/j.expneurol.2015.02.023
- [60] Gerber YN, Sabourin JC, Rabano M, Vivanco M, Perrin FE. Early functional deficit and microglial disturbances in a mouse model of amyotrophic lateral sclerosis. *PLoS One* (2012) 7(4):e36000. doi:10.1371/journal.pone.0036000
- [61] Wu DC, Re DB, Nagai M, Ischiropoulos H, Przedborski S. The inflammatory NADPH oxidase enzyme modulates motor neuron degeneration in amyotrophic lateral sclerosis mice. *Proc Natl Acad Sci U S A* (2006) 103:12132-7. doi:10.1073/pnas.0603670103
- [62] Boillee S, Vande Velde C, Cleveland DW. ALS: a disease of motor neurons and their nonneuronal neighbors. *Neuron* (2006) 52:39-59. doi:10.1016/j.neuron.2006.09.018
- [63] Henn A, Lund S, Hedtjarn M, Schratzenholz A, Porzgen P, Leist M. The suitability of BV2 cells as alternative model system for primary microglia cultures or for animal experiments examining brain inflammation. *ALTEX* (2009) 26:83-94. doi:10.14573/altex.2009.2.83
- [64] Horvath RJ, Natile-McMenemy N, Alkaitis MS, Deleo JA. Differential migration, LPS-induced cytokine, chemokine, and NO expression in immortalized BV-2 and HAPI cell lines and primary microglial cultures. *J Neurochem* (2008) 107:557-69. doi:10.1111/j.1471-4159.2008.05633.x
- [65] Olmos G, Llado J. Tumor necrosis factor alpha: a link between neuroinflammation and excitotoxicity. *Mediators Inflamm* (2014) 2014:861231. doi:10.1155/2014/861231
- [66] Obal I, Jakab JS, Siklos L, Engelhardt JI. Recruitment of activated microglia cells in the spinal cord of mice by ALS IgG. *Neuroreport* (2001) 12(11):2449-52. doi:10.1097/00001756-200108080-00032

- [67] Bozic I, Savic D, Laketa D, Bjelobaba I, Milenkovic I, Pekovic S, et al. Benfotiamine attenuates inflammatory response in LPS stimulated BV-2 microglia. *PLoS One* (2015) 10(2):e0118372. doi:10.1371/journal.pone.0118372
- [68] Zhao W, Beers DR, Henkel JS, Zhang W, Urushitani M, Julien JP, et al. Extracellular mutant SOD1 induces microglial-mediated motoneuron injury. *Glia* (2010) 58(2):231-43. doi:10.1002/glia.20919
- [69] Lewis KE, Rasmussen AL, Bennett W, King A, West AK, Chung RS, et al. Microglia and motor neurons during disease progression in the SOD1G93A mouse model of amyotrophic lateral sclerosis: changes in arginase1 and inducible nitric oxide synthase. *J Neuroinflammation* (2014) 11:55. doi:10.1186/1742-2094-11-55
- [70] Beckman JS, Koppenol WH. Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and ugly. *Am J Physiol* (1996) 271:C1424-37
- [71] Zhao W, Xie W, Le W, Beers DR, He Y, Henkel JS, et al. Activated microglia initiate motor neuron injury by a nitric oxide and glutamate-mediated mechanism. *J Neuropathol Exp Neurol* (2004) 63:964-77. doi:10.1093/jnen/63.9.964
- [72] Kaneko YS, Ota A, Nakashima A, Mori K, Nagatsu I, Nagatsu T. Regulation of oxidative stress in long-lived lipopolysaccharide-activated microglia. *Clin Exp Pharmacol Physiol* (2012) 39:599-607. doi:10.1111/j.1440-1681.2012.05716.x
- [73] Chatterjee S, Noack H, Possel H, Wolf G. Induction of nitric oxide synthesis lowers intracellular glutathione in microglia of primary glial cultures. *Glia* (2000) 29:98-101. doi:10.1002/(SICI)1098-1136(20000101)29:1<98::AID-GLIA10>3.0.CO;2-B
- [74] Schulz JB, Lindenau J, Seyfried J, Dichgans J. Glutathione, oxidative stress and neurodegeneration. *Eur J Biochem* (2000) 267:4904-11. doi:10.1046/j.1432-1327.2000.01595.x
- [75] Przedborski S, Donaldson D, Jakowec M, Kish SJ, Guttman M, Rosoklija G, et al. Brain superoxide dismutase, catalase, and glutathione peroxidase activities in amyotrophic lateral sclerosis. *Ann Neurol* (1996) 39:158-65. doi:10.1002/ana.410390204
- [76] Fujita K, Yamauchi M, Shibayama K, Ando M, Honda M, Nagata Y. Decreased cytochrome c oxidase activity but unchanged superoxide dismutase and glutathione peroxidase activities in the spinal cords of patients with amyotrophic lateral sclerosis. *J Neurosci Res* (1996) 45:276-81. doi:10.1002/(SICI)1097-4547(19960801)45:3<276::AID-JNR9>3.0.CO;2-A
- [77] Shaw IC, Fitzmaurice PS, Mitchell JD, Lynch PG. Studies on cellular free radical protection mechanisms in the anterior horn from patients with amyotrophic lateral sclerosis. *Neurodegeneration* (1995) 4:391-6. doi:10.1006/neur.1995.0047
- [78] Harguindeguy S, Reshkin SJ, Orive G, Arranz JL, Anitua E. Growth and trophic factors, pH and the Na<sup>+</sup>/H<sup>+</sup> exchanger in Alzheimer's disease, other neurodegenerative diseases and cancer: new therapeutic possibilities and potential dangers. *Curr Alzheimer Res* (2007) 4:53-65. doi:10.2174/156720507779939841